

Hasil Review Artikel/manuskrip pada Jurnal Veterinar World

Judul ; *Shigella flexneri* vaccine development: Oral administration of peptides derived from the 49.8 kDa pili protein subunit activates the intestinal immune response in mice

Penulis ; Khoirul Anam, Agustina Tri Endharti, Sri Poeranto, Hidayat Sujuti, Dwi Yuni Nur Hidayati, Sumarno Reto Prawiro,

Veterinary World

For your questions send email to editor@veterinaryworld@gmail.com

Status of my Manuscripts waiting for Editor's first View

Mms No	Submission Date	Title	Authors	Status
VETWORLD-2021-09-535	2021-09-25	Initial Version (DOC): Initial Full Text (.doc) Initial Version (PDF): The Study of Epitope of 49.8 kDa Pili Protein Subunit <i>Shigella flexneri</i> as Prospective Shigellosis Vaccine to Improve Mucosal Immune Response in Mice Through Oral Immunization	Khoirul Anam, Khoirul Anam, Agustina Tri Endharti, Sri Poeranto, Hidayat Sujuti, Dwi Yuni Nur Hidayati, Sumarno Reto Prawiro,	Waiting for Editor's View

Veterinary World

For your questions send email to editor@veterinaryworld@gmail.com

Status of my Manuscripts under Review

Mms No	Submission Date	Title	Authors	Status
VETWORLD-2021-09-535	2021-09-25	Initial Version (DOC): Initial Full Text (.doc) Initial Version (PDF): The Study of Epitope of 49.8 kDa Pili Protein Subunit <i>Shigella flexneri</i> as Prospective Shigellosis Vaccine to Improve Mucosal Immune Response in Mice Through Oral Immunization	Khoirul Anam, Khoirul Anam, Agustina Tri Endharti, Sri Poeranto, Hidayat Sujuti, Dwi Yuni Nur Hidayati, Sumarno Reto Prawiro,	With Editorial Board

Browser tabs: (9.877 belum dibaca) x | Google x | WhatsApp x | Pengumuman Pendid... x | Veterinary World - C... x | Veterinary World : e... x

Address bar: <https://www.ejmanager.com/my/vetworld/submit.php?isi=ur&lng=>

Veterinary World

Navigation: Home/About | Guide for Authors | Search Journal | Submit New Manuscript | Status of my Articles | Personal Information | LOGOUT (Khoiril Anam) | eManager 3.7

For your questions send email to editor@veterinaryworld@gmail.com | Personal Messages(0)

Status of my Manuscripts under Review

Mas No	Submission Date	Title	Authors	Status
VETWORLD-2021-09-535	2021-09-25	Initial Version (DOC): Initial Full Text (.doc) Initial Version (PDF): The Study of Epitope of 49.8 kDa Pili Protein Subunit Shigella flexneri as Prospective Shigellosis Vaccine to Improve Mucosal Immune Response in Mice Through Oral Immunization	Khoiril Anam, Khoiril Anam, Khoiril Anam, Agustina Tri Endharti, Siti Poeranto, Hidayat Sujati, Dwi Yuni Nur Hidayati, Sumarno Reto Prawiro,	Under Review

Windows taskbar: Type here to search | 33°C | 11:02 03/11/2021

Browser tabs: What's x | Surat x | (9.905 x | 2-163: x | Goog x | Veteri x | Veteri x | VWJ: x | acader x | Devel: x | Veteri x

Address bar: <https://mail.yahoo.com/d/folders/1/messages/137182.intl=id&lang=id-ID&partner=none&src=fp>

yahoo!mail

Temukan pesan, dokumen, foto, atau orang

Tulis | Kembali | Arsipkan | Pindahkan | Hapus | Spam

Email Mas... 999+

Belum Dibaca | Berbingung | Draft | Terkirim | Arsip | Spam | Sampah | Lebih sedikit | Tampilan | Sembunyikan | Folder Baru | tidak berjudul... | tidak berjudul

Revised Article Submission

Veterinary World <noreply@ejmanager.com>
Kepada: sanambwi@yahoo.co.id | Sab, 25 Des jam 17.37

Dear Khoiril Anam,

Your REVISED ARTICLE entitled **The Improvement of Intestinal Immune Response Following Oral Immunization of Epitope of 49.8 kDa Pili Protein Subunit Shigella flexneri in Mice Model Revealed Potential Vaccine Candidate Development** (Mns No VETWORLD-2021-09-535) has been received by Veterinary World.


You could follow status of your manuscript by login to your author account at www.ejmanager.com

Thank you for submitting your REVISED version of your article.

Best regards,
Editor
Veterinary World
<http://www.veterinaryworld.org>

JOURNAL CONTACT EMAIL: editor@veterinaryworld@gmail.com

Windows taskbar: Type here to search | 27°C



Technical/Copyediting by Sinjore – 19/01/2022

RESEARCH ARTICLE

***Shigella flexneri* vaccine development: Oral administration of peptides derived from the 49.8 kDa pili protein subunit activates the intestinal immune response in mice**

Khoiril Anam^{1,2} {ORCID:0000-0001-7676-2827}, Agustina Tri Endharti³ {ORCID:0000-0002-2062-5740}, Sri Poeranto³ {ORCID:0000-0002-2366-4657}, Hidayat Sujuti⁴ {ORCID:0000-0001-6568-3584}, Dwi Yuni Nur Hidayati⁵ {ORCID:0000-0001-8116-4573} and Sumarno Reto Prawiro⁵ {ORCID:0000-0002-3718-4806}

1. Doctoral Program in Medical Science, Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia; 2. Medical Laboratory Technology of Study Program Institute of Health and Science Technology Wiyata Husada, Samarinda, Indonesia; 3. Department of Parasitology, Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia; 4. Department of Biochemistry, Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia; 5. Department of Clinical Microbiology, Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia.

Corresponding author: Sumarno Reto Prawiro, e-mail: retoprawiros@yahoo.com

Co-authors: KA: sanambwi@yahoo.co.id, ATE: tinapermana@yahoo.com, SP: dr.poer@gmail.com, HS: hidayatsujuti@yahoo.com, DYNH: yuninur83@yahoo.com

Received: 27-09-2021, **Accepted:** 05-01-2022, **Published online:** ***

doi: *** **How to cite this article:** Anam K, Endharti AT, Poeranto S, Sujuti H, Hidayati DYN, Prawiro SR (2022) *Shigella flexneri* vaccine development: Oral administration of peptides derived from the 49.8 kDa pili protein subunit activates the intestinal immune response in mice, *Veterinary World*, 15(2): 0-0

Abstract

Background and Aim: The morbidity and mortality of *Shigella* infections remain a global challenge. Epitope-based vaccine development is an emerging strategy to prevent bacterial invasion. This study aimed to identify the ability of the 49.8 kDa pili subunit adhesin protein epitope of *Shigella flexneri* to induce an intestinal immune response in mice.

Materials and Methods: Thirty adult male Balb/c mice were divided into a control group, cholera toxin B subunit (CTB) group, CTB+QSSTGTNSQSDLDS (pep_1) group, CTB+DTTITKAETKTVTKNQVVDTPVTDAAK (pep_2) group, and CTB+ATLGATLNRLDFNVNKK (pep_3). We performed immunization by orally administering 50 µg of antigen and 50 µl of adjuvant once a week over 4 weeks. We assessed the cellular immune response by quantifying T helper 2 (Th2) and Th17 using flow cytometry. In addition, we assessed the humoral immune response by quantifying interleukin (IL-4), IL-17, soluble immunoglobulin A (sIgA), and β-defensin using enzyme-linked immunoassay. Statistical analysis was performed using one-way analysis of variance and Kruskal–Wallis test.

Results: Peptide oral immunization increases the cellular immune response as reflected by

Commented [Editor1]: Remark:
These group names are not very informative. In the main text, it is good to have short group names but in the abstract, consider being more descriptive. It would be helpful for the readers to know what pep_1, pep_2 and pep_3 are.

We have edited it. Please check.

Commented [Microsoft2R1]: We did check it. Thank you

the increase of Th2 (p=0.019) and Th17 (p=0.004) cell counts, particularly in the CTB_pep_1 group. Humoral immune response activation was demonstrated by the increase in IL-4 levels, especially in the CTB+pep_3 group (p=0.000). The IL-17 level **rise** significantly in the CTB+pep_1 group (p=0.042). The mucosal immune response was demonstrated by the sIgA levels increase in the CTB+pep_3 group (p=0.042) and the β -defensin protein levels (p=0.000).

Commented [Microsoft3]: We change into "rise "

Conclusion: All selected peptides activated the cellular and humoral immune responses in the intestine of mice. Further studies are necessary to optimize antigen delivery and evaluate whether the neutralizing properties of these peptides allow them to prevent bacterial infection.

Keywords: epitope, immune response, oral, *Shigella flexneri*, shigellosis.

<H1>Introduction

Shigellosis is an acute gastrointestinal infectious disease characterized by bloody stool diarrhea. Its mortality and morbidity rates are particularly high in low- and middle-income economy countries. It is highly prevalent in children under 5 years, particularly in poor sanitary conditions and overcrowded populations [1]. One of the pathogenic causes of

shigellosis is *Shigella flexneri* species. This species is frequently recorded in shigellosis cases worldwide [2-4]. The disease burden and antimicrobial resistance of *S. flexneri* infection remain global challenges and overcoming them require developing new strategies [5,6].

Shigella invasion is initiated by bacterial adhesion to the enteric epithelial lining. Therefore, the pili portion of *Shigella* is crucial for bacterial host infection. Sumarno *et al.* [7] reported that the pili of *Shigella dysenteriae* contain a hemagglutinin protein weighing 49.8 kDa that acts as an adhesin [8]. Other *Shigella* species such as *S. flexneri*, *Shigella sonnei*, and *Shigella boydii* have similar pili proteins [9]. The importance of pili proteins was demonstrated by administering the 37.7 kDa pili protein subunit *Vibrio cholerae* 01, which stimulated the mucosal immune response by increasing soluble immunoglobulin A (sIgA) levels [10].

Orally administering the 49.8 kDa pili protein subunit of *S. dysenteriae* yielded a similar response in mice [11].

T helper 2 (Th2) and Th17 cells promote the mucosal immune response toward *Shigella* infection. The Th2 cells mediate the humoral immune response by secreting interleukin 4 (IL-4), IL-5, IL-6, and IL-13 [12]. The immunoglobulin A (IgA) immune response induced by Th2 cells inhibits the bacterial attachment to epithelial cells [13]. Moreover, Th17 cells have

Commented [Editor4]: Remark:
Please add the species to which this subunit was administered in that experiment ("to mice" or "to rats" for example).

Commented [Microsoft5R4]: We provide "in mice" as additional

a key role in the pathogenesis of autoimmune and inflammatory diseases [14]. The differentiated Th17 cells secrete the cytokines IL-17 and IL-22. These cytokines promote the release of secretory IgA (sIgA) in the intestinal lumen and contribute to antimicrobial proteins secretion [15,16]. Antimicrobial peptides (AMPs) also play crucial roles in the adaptive immune system and pathogen elimination [17].

Despite the role of pili proteins in humoral immune response induction, *S. flexneri* immunization combined with cholera toxin B subunit (CTB) and Sumbawa horse milk administration increases sIgA and β -defensin levels [18]. The developing countries would greatly benefit from efficacious and safe vaccines against *Shigella*. However, such vaccines require further development [3]. The current vaccine candidates are limited to attenuated strain vaccines, conjugate vaccines, subunit vaccines (Invaplex, OMP, T3SS), and combination vaccines (*Shigella*-ETEC and *Shigella*-Salmonella) [19]. The current efforts toward vaccines or early detection diagnostic tests using pili proteins are promising [20]. Identifying the adhesin protein sequences revealed protein epitopes potentially useful for vaccine development. Pore *et al.* [21] analyzed the amino acid sequence of the 34 kDa outer membrane protein (OmpA) of *S. flexneri*. Similarly, Sharma *et al.* [22] performed epitope prediction modeling on the outer membrane proteins (OMPs) of *S. flexneri* 2a.

Commented [AS6]:
Reference no.3 and 19 were same so, we have removed duplicate reference no.19 and rearranged reference no. in the text as well as in the reference section.

Commented [Microsoft7R6]: Thank you

The chemical stability, easy production, and minimal infection potential of peptides make them attractive vaccine candidates. The previous peptide vaccines have been developed and approach the clinical stage. Several studies demonstrated the effectiveness of epitope-based vaccine design in stimulating long-term antibody production and protective immunity against *V. cholera* [23-25]. In addition, our previous study demonstrated the epitope sequence of the 49.8 kDa pili protein from *S. flexneri*-induced anti-peptide serum antibodies and prevented leaking in the enterocyte of mice [26].

This study aimed to document the antigenic properties of the epitopes of the 49.8 kDa *S. flexneri* pili protein by assessing their ability to induce cellular and humoral immune responses in the intestine of mice. This study should provide fundamental data for the development of epitope-based vaccines against *S. flexneri* as part of the effort to overcome the shigellosis infection issues.

<H1>Materials and Methods

<H2>Ethical approval

Adult male Balb/c mice were obtained and housed in Experimental Animal Laboratory, Faculty of Medicine, Universitas Brawijaya. Animals were kept under maintained of

humidity, temperature, and regular dark light cycle. Animals had free access to drink as well as standard laboratory food [27]. All procedures were previously approved by Research Ethic Committee, Universitas Brawijaya (no.1192-KEP-UB).

<H2>Study period and location

This study was performed from January to March 2021 at Experimental Animal Laboratory, Clinical Parasitology Laboratory, and Biomedical Central Laboratory, Faculty of Medicine, Universitas Brawijaya

<H2>Experimental design

We separated 30 adult male Balb/C mice (6-8 weeks old) into control group, CTB group, CTB+QSSTGTNSQSDLDS (pep_1) group, CTB+DTTITKAETKTVTKNQVVDTPVTDAAK (pep_2) group, and CTB+ATLGATLNRLDFNVNKK (pep_3). We used CTB as an adjuvant. We performed immunization by orally administering 50 µg of peptide in 50 µl of adjuvant once a week for 4 weeks. We sacrificed the mice 1 week after the last immunization [28]. As antigens, we used synthetic peptides from the 49.8 kDa *S. flexneri* pili protein named pep_1, pep_2, and pep_3

(Apical Scientific Sdn. Bhd., Malaysia).

<H2>Peptide sequencing and modeling

Isolated protein from *Shigella flexneri* philli was denatured in 100°C of Laemmli buffer and electrophoresed in SDS-PAGE on 4% gel. Coomassie brilliant blue was applied to visualize protein bands. The protein band at 49,8 kDa was excised and dried for sequencing analysis at Proteomic International (Australia) according to in-gel digestion method and spectral analysis using an electrospray ionization mass spectrometer (Agilent 1260 Infinity HPLC system, USA) paired with an Agilent 6540 mass spectrometer. Data was analyzed using Mascot software (Matrix Science, USA) with Ludwig NR database [29]. We modeled the secondary protein structure of the peptides using the PEP-FOLD3 server (<http://mobyli.rpbs.univ-paris-diderot.fr/cgi-bin/portal.py#forms::PEP-FOLD3>).

<H2>Molecular analysis

<H3>Flow cytometry

We counted Th2 and Th17 cells in intestinal mesenteric tissue using a flow cytometry apparatus (BD FACS Melody, USA). We performed Th2 and Th17 double staining using

Commented [AS8]:
Company and country name for this product.

FITC-labeled anti-mouse CD4 antibody (BioLegend, San Diego, CA, USA) and PerCP-labeled anti-mouse IL-4 for Th2 (BioLegend, San Diego, CA, USA) and FITC-labeled anti-mouse CD4 antibody and PE-labeled anti-mouse IL-17 (BioLegend, San Diego, CA, USA) for Th17 [30].

Commented [AS9]:

Company and country name for this product.

Commented [AS10]:

Company and country name for this product.

<H3>Enzyme-linked immunoassay

We quantified IL-4 and IL-17 from intestinal mesentery and sIgA and β -defensin from intestinal mucosal tissue using an enzyme-linked immunoassay (Zenix-320 microplate reader) according to the manufacturer's instructions (Bioassay Technology Laboratory, Shanghai, China) [28].

<H2>Statistical analysis

Data were presented as mean \pm standard deviation (SD). The differences between groups were analyzed by one-way analysis of variance and Kruskal–Wallis test using SPSS software ver. 23 (IBM, USA). The significant value was set at $p < 0.05$.

Commented [TPS11]: Kindly provide manufacturer details

<H1>Results

Figure-1 shows the secondary structures of the three selected peptides obtained by modeling.

Pep_1, Pep_2, and Pep_3, respectively, contain a helix, a coil (thread), and a sheet. Figure-2 shows the cellular immune responses. The immunization groups had significantly higher Th2 and Th17 levels than the control group. The CTB+pep_1 group had the highest Th2 (p=0.019) and Th17 levels (p=0.004). Regarding the humoral immune response, the immunization groups, particularly the CTB+pep_3 group (p=0.000), had increased IL-4 levels. However, only the CTB+pep_1 group (p=0.042) had a significantly higher IL-17 level than the control group (Figure-3). Finally, Figure-4 presents the mucosal immune response results. The CTB+pep_3 group had a significantly elevated sIgA level (p=0.042). Moreover, the immunization groups had higher levels of the natural immune response marker β -defensin (p=0.000).

<H1>Discussion

S. flexneri enters the epithelial cells by transiting through M cells and encounters resident macrophages. Pro-inflammatory signals released by macrophages and epithelial cells then activate a natural immune response involving natural killer cells and polymorphonuclear cells [31]. *Shigella* relies on the production of effector proteins that contribute to the manipulation of the infectious process as a specific mechanism to invade host cells [32]. Two types of

Commented [Editor12]: Remark:
Please confirm that this corresponds to what you meant.

Commented [Microsoft13R12]: yes It is corresponds to our meant

attachment proteins of *Shigella* are fimbriae/pili proteins and the OMPs. Pili proteins and OMPs are virulence factors for bacterial colonization that causes infection [33,34]. Therefore, developing OMP epitope-based vaccines against *Shigella* infection are a sensible strategy [22]. We identified the potential of *S. dysenteriae*, *S. flexneri*, *S. sonnei*, and *S. boydii* pili proteins as adhesive proteins that can be used for *Shigellosis* vaccine development [7,9]. The current study revealed the development of an epitope-based vaccine candidate against *Shigella* with good prospects and safety. We utilized three antigenic peptides derived from the 49.8 kDa pili protein subunit of *S. flexneri* and demonstrated that they induced the immune cellular and humoral responses as protection mechanisms against *Shigella* [35].

Oral vaccination against shigellosis can increase local immune response as linear with the pathogenesis of *Shigella*. In this study, we administered immunizing compounds orally and assessed the mucosal immune response through sIgA and β -defensin, which are essential immune system proteins in enteric infections. Preventing enteric mucosal infections in humans require IgA secretion. *S. flexneri*-infected patients had significantly more cells secreting circulating antibodies than healthy individual [36]. Activated CD4+ T cells differentiate into Th1, Th2, Th17, and regulatory T cells [37,38]. Th2 cells produce mediator cytokines such as IL-4, IL-5, and IL-10, which stimulate the production of plasma cell

Commented [Editor14]: Remark:
For clarity, consider adding who these patients are compared to (“... than healthy subjects” or “... than patients infected with other bacteria” for example).

antibodies, including IgA. An intensive increase in IgA production reflects the stimulation of the mucosal defense [39]. Activated Th17 cells differentiate and secrete IL-17 and IL-22. IL-17 increases the secretion of sIgA into the lumen [16]. In addition, IL-17 and IL-22 can help intestinal epithelial cells produce antimicrobial proteins [15]. IL-17 and IL-22 strongly stimulate AMPs secretion by epithelial cells [40].

Based on the mechanism of *Shigella* pathogenesis and immune response, we measured the ability of our peptides to stimulate the immune responses by quantifying Th2, IL-4, and sIgA. Treating mice with 49.8 kDa *S. flexneri* pili protein subunit-based peptides increased the Th2 cellular immune response. The three peptides tested (pep_1, pep_2, and pep_3) were immunogenic. The ability of these peptides to induce IL-4 production by Th2 cells is encouraging for vaccine design [41]. This correlates with the previous studies that reported increased lymphocyte proliferation after vaccination with the rIpaB protein of *S. flexneri* and rGroEL of *Salmonella typhi* [42]. *Salmonella* secretes effector I (SseI or SrfH) and the flagellin protein FliC, which activate T cells. Inducing the immune response of CD4 T-cells using the epitopes of these proteins protects against infection [43,44]. The proliferation of Th2 cells increases the levels of IL-4, a cytokine that plays a key role in numerous biological activities [45].

Commented [Editor15]: Remark:
Please confirm that this corresponds to what you meant.

Commented [Microsoft16R15]: yes It is corresponds to our meant

The cytokine IL-4 is part of the humoral immune response. The CTB+pep_2 and CTB+pep_3 groups had higher IL-4 levels than the control group. A previous study reported a similar IL-4 production after administering a peptide derived from the EpiMix protein of *S. flexneri* [46]. Another study revealed that the recombinant chimeric protein rMESF of *S. flexneri* significantly increased IL-4 levels through all tested application routes, that is, intranasal, intravenous, intramuscular, and subcutaneous [47].

Mucosal antibody production is proportional to Th2 and IL-4 levels. Th2 cells produce IL-4

cells, activating sIgA production, which facilitate B cells activation and antibody release. B

cells in the lamina propria produce IgA, protecting the mucosal tissue and neutralizing

microbes before invading the mucosal lumen. sIgA acts the outer bacterial membrane and

inhibits the bacterial attachment to mucosal surfaces [36,48]. The increase in sIgA levels in

this study suggested that our vaccine candidates stimulated the mucosal immune responses.

The vaccine using *S. flexneri* rMESF multiepitope protein increases sIgA-specific antibodies

both in the mucus layer and system wide [47]. EpiMix, *S. flexneri* protein fragment, produces

a specific sIgA mucosal immune response from immunized animal fecal samples [46]. The

presence of pathogen-specific sIgA is important to prevent the infection of mucosal tissues.

IgA is an immunoglobulin essential *Shigella* infection and reinfection prevention and

Commented [Editor17]: Remark:
Please confirm that this corresponds to what you meant.

Commented [Microsoft18R17]: yes It is corresponds to
our meant

treatment. Therefore, IgA production inducing antibodies are the primary protection mechanism against *Shigella* infection [49].

Our study assessed the immune response through Th17 cells, IL-17, and β -defensin. Th17 cells produce the main cytokines, namely, IL-17 and IL-22. In addition, Th17 cells stimulate the production of antimicrobial substances called defensins that function like locally produced endogenous antibiotics [38]. Our peptides successfully activated the Th17 cellular immune response. The ability of these peptides to activate Th17 cells demonstrates their strong potential as shigellosis vaccine candidates. Numerous scientific reports show that Th17 cells and IL-17 can induce the production of antimicrobial cytokines and proteins that contribute to the host's defense system against intestinal pathogens. IL-17 increases sIgA secretion into the lumen as well [16]. *Shigella* infection induces IL-17 and IL-22 by Th17 cells, increasing the host's defense against bacteria and fungi [15].

IL-17 is a cytokine mainly produced by Th17 cells involved in the protection against pathogens in mucosal tissues [50,51]. Therefore, an increase in IL-17 levels indicates the efficacy of a vaccine [52], and pep_1, pep_2, and pep_3 increased the IL-17 levels. Our results are consistent with the previous studies [47]. IL-17 and IL-22 can stimulate the

production of antimicrobial proteins by intestinal epithelial cells [15]. Antimicrobial proteins are a gastrointestinal mucosal defense against pathogens [53].

AMPs are part of the natural humoral immune systems of the innate immune response against various pathogenic organisms. They govern the regulatory signals for the innate and adaptive immune systems to eliminate pathogens [17]. Most mammals express AMPs from the defensin and cathelicidin families [54]. Defensins play an important role in biological processes because of their antimicrobial and immunomodulatory activities [17]. β -defensin has a positive charge making it effective against negatively charged bacteria [55]. Our results showed that our peptides increased β -defensin levels. The presence of β -defensins is crucial because they can attract inflammatory cells such as neutrophils, T cells, and macrophages [56].

Despite the high efficacy of peptide-based oral vaccines, the utilization of peptides as antigens for oral vaccine development faces several limitations, such as proteolysis degradation. However, using delivery systems, coadministering protease inhibitors can enhance the effect of vaccine candidates [57,58]. Therefore, we aim to enhance the efficacy of our peptide candidate and optimize antigen delivery systems in further studies.

<H1>Conclusion

It can be concluded that peptides derived from the 49.8 kDa *S. flexneri* pili protein subunit as prospective shigellosis vaccines by demonstrating their ability to induce the mucosal immune response. The pep_3 peptide sequence was particularly efficient. Further studies involving peptide delivery systems and neutralization examination through bacterial colony growth inhibition are necessary to realize the potential of the selected peptide.

<H1>Authors' Contributions

SRP and HS were conceptualized the research; KA, ATE and DYNH were designed the research and data acquisition; KA and SRP were performed data analysis, data interpretation and drafting the manuscript; All author were revised manuscript for critically content as well as final approval of the version to be published; SRP was provided agreement to be accountable.

<H1>Acknowledgments

The authors would like to thank Wahyuda Ngatiril Lady from Central Laboratory of Biomedics and Heni Indrawati from Clinical Parasitology Laboratory, Faculty of Medicine,

Commented [TP519]: Kindly provide the text part as per the journal style.

Commented [Microsoft20R19]: we just provide the additional

Commented [TP521]: Kindly provide Grant number

Commented [Microsoft22R21]: We did not

Universitas Brawijaya, Malang, Indonesia, for technical assistance.

<H1>Competing Interests

The authors declare that they have no competing interests.

<H1>Publisher's Note

Veterinary World remains neutral with regard to jurisdictional claims in published institutional affiliation.

<H1>References

1. Niyogi, S.K. (2005) Shigellosis. *J. Microbiol.*, 43(2): 133-143.
2. Al-Moyed, K.A., Harmal, N.S., Al-Harasy, A.H. and Al-Shamahy, H.A. (2006) Increasing single and multi-antibiotic resistance in *Shigella* species isolated from shigellosis patients in Sana'a, Yemen. *Saudi Med. J.*, 27(8): 1157-1160.
3. Mani, S., Wierzba, T. and Walker, R.I. (2016) Status of vaccine research and development for *Shigella*. *Vaccine*, 34(26): 2887-2894.

4. Nikfar, R., Shamsizadeh, A., Darbor, M., Khaghani, S. and Moghaddam, M. (2017) A Study of prevalence of *Shigella* species and antimicrobial resistance patterns in paediatric medical center, Ahvaz, Iran. *Iran. J. Microbiol.*, 9(5): 277-282.
5. Gaurav, A., Singh, S.P., Gill, J.P.S., Kumar, R. and Kumar, D. (2013) Isolation and identification of *Shigella* spp. from human fecal samples collected from Pantnagar, India. *Vet. World*, 6(7): 376-379.
6. Khaghani, S., Shamsizadeh, A., Nikfar, R. and Hesami, A. (2014) *Shigella flexneri*: A three-year antimicrobial resistance monitoring of isolates in a Children Hospital, Ahvaz, Iran. *Iran. J. Microbiol.*, 6(4): 225-229.
7. Sumarno, R.P., Avanita, A.S., Winarsih, S., Hidayat, S. and Nurhidayati, D.Y. (2015) Haemagglutination of *Shigella dysenteriae* subunit pili protein with and anti-haemagglutination of *Shigella dysenteriae* subunit pili protein as a molecule adhesin in mouse enterocytes. *Afr. J. Med. Res.*, 9(11): 781-787.
8. Agustina, W., Fitri, L.E., Raras, T.Y.R., Siswanto, B. and Prawiro, S.R. (2012) Antibody protein hemagglutinin subunit pili with MW 49,8 kDa *Shigella dysenteriae* can inhibit *Shigella dysenteriae* adhesin on mice enterocyte. *IOSR J. Pharm.*, 2(5): 13-20.

9. Anam, K., Nurdiana., Herowati, T.E., Sujuti, H. and Sumarno, R.P. (2016) Cross immunity among pili sub-unit Hemagglutinin and pili sub-unit anti hemagglutinin proteins of *Shigella* spp. *Int. J. Pharm Pharm. Res. Human*, 7(2): 19-30.
10. Faisal, R., Sumarno. and Handono, K. (2010) Sumbawa fermented horse milk as immunostimulants for 37.8 kDa *V. cholerae* vaccine. *J. Braw Med.*, 26(1): 225-230.
11. Setyorini, A., Utami, Y.D., Wijayanto, E., Winarsih, S., Noorhamdani, A.S. and Prawiro, S.R. (2013) Protective of adhesin molecules pili 49,8 kDa *Shigella dysenteriae* conjugated with ISCOM against bacterial colonization and colonic epithelial cells damage in mice. *Int. J. Trop. Med.*, 8(1): 19-26.
12. Yang, Y., Liu, K., Chen, Y., Gong, Y. and Liang, Y. (2019) Indoleamine 2,3-dioxygenase (IDO) regulates Th17/Treg immunity in experimental IgA nephropathy. *Folia Biol. (Praha)*, 65(2): 101-108.
13. Yanagita, M., Hiroi, T., Kitagaki, N., Hamada, S., Ito, H., Shimauchi, H., Murakami, S., Okada, H. and Kiyono, H. (1999) Nasopharyngeal-associated lymphoreticular tissue (NALT) immunity: Fimbriae-specific Th1 and Th2 cell-regulated IgA responses for the inhibition of bacterial attachment to epithelial cells and subsequent inflammatory cytokine production. *J. Immunol.*, 162(6): 3559-3565.

14. Dong, C. (2008) Th17 cells in development: An updated view of their molecular identity and genetic programming. *Nat. Rev. Immunol.*, 8(5): 337-348.
15. Sellge, G., Magalhaes, J.G., Konradt, C., Fritz, J.H., Salgado-Pabon, W., Eberl, G., Bandeira, A., Di Santo, J.P., Sansonetti, P.J. and Phalipon, A. (2010) Th17 cells are the dominant T Cell subtype primed by *S. flexneri* mediating protective immunity. *J. Immunol.*, 184(4): 2076-2085.
16. Cao, A.T., Suxio, Y., Bin, G., Co, E. and Yingzi, C. (2012) Th17 Cells Upregulate Polymeric Ig receptor and intestinal homeostasis. *J. Immunol.*, 189(9): 4666-4673.
17. Shin, D.M. and Kyeong Jo, E. (2011) Antimicrobial peptides in innate immunity against mycobacteria. *Immune. Network*, 11(5): 245-252.
18. Utami, W.J., Wiwik, A., Winarsih, S., Sujuti, H. and Sumarno. (2017) The fermented horse milk can as an immune stimulant pili sub units of *S. flexneri* protein conjugated with CTB in mice. *Acta Sci. Pharm. Sci.*, 1(2017): 30-37.
19. Walker, R.I., Wierzba, T.F., Mani, S. and Bourgeois, A.L (2017) Vaccines against *Shigella* and enterotoxigenic *Escherichia coli*: A summary of the 2016 VASE conference. *Vaccine*, 35(49 Pt A): 6775-6782.

20. Suharsono, H., Muttaqin, Z., Tenaya, I.W.M., Agustina, K.K. and Prawiro, S.R. (2019) Antigen of 49.6-kDa subunit pili protein of *Helicobacter pylori* as a potential biomarker for early and rapid detection of H. pylori infection. *Vet. World*, 12(6): 769-773.
21. Pore, D., Mahata, N., Pal, A. and Chakrabarti, M.K. (2011) Outer membrane protein A (OmpA) of *Shigella flexneri* 2a, induces protective immune response in a mouse model. *PLoS One*, 6(7): e22663.
22. Sharma, D., Patel, S., Padh, H. and Desai, P. (2016) Immunoinformatic identification of potential epitopes against shigellosis. *Int. J. Pept. Res. Ther.*, 22(4): 481-495.
23. Patronov, A. and Doytchinova, I. (2013) T-cell epitope vaccine design by immunoinformatics. *Open Biol.*, 3(1): 120139.
24. Nezafata, N., Karimia, Z., Eslamia, M., Mohkama, M., Zandiana, S. and Ghasemia, Y. (2016) Designing an efficient multiepitope peptide vaccine against *Vibrio cholerae* via combined immunoinformatics and protein interaction based approaches. *Comput. Biol. Chem.*, 62(June): 82-95.

25. Ghazi, F.M.P. and Gargari, S.L.M. (2017) Synthetic peptides mimicking lipopolysaccharide as a potential vaccine candidates against *Vibrio cholerae* serogroup O1. *Iran. J. Microbiol.*, 9(4): 244-250.
26. Sumarno., Anam, K., Utami, Y.W., Nurhidayati, Y., Winarsih, S. and Sujuti, H. (2019) The response of anti-peptide serum antibodies ATLGATLNRLDFNVNKK derived from *S. flexneri* protein sub-unit pili toward activity the hemagglutination and protection from leaking solution enterocyte in mice. *Arch. Immunol. Allergy*, 2(2): 49-57.
27. Kurnianingsih, N., Ratnawati, R., Nazwar, T.A., Ali, M. and Fatchiyah, F. (2020) The behavioral effect of anthocyanin from purple sweet potatoes on prenatally stressed offspring mice. *Syst. Rev. Pharm.*, 11(10): 482-490.
28. Guli, M., Illiandri, O., Winarsih, S., Barlianto, W. and Prawiro, S.R. (2021) Immunoenhancing effect of *Lactobacillus reuteri* on immunized mice intestine using cholerae toxin subtype B. *J. Adv. Pharm. Edu. Res.*, 11(1): 11-17.
29. Bringans, S., Soren, E., Tulene, K., Gopalakrishnakone, P., Andreja, L., Robert, L. and Lipscombe, R. (2008) Proteomic analysis of the venom of *Heterometrus longimanus* (Asian black scorpion) Scott. *Proteomics*, 8(5): 1081-1096.

30. Prawiro, S.R., Anam, K., Prabowo, B., Bramanthi, R., Fitriainingsih, A.A., Hidayati, .
D.Y.N., Imawati, S., Fitria, E. and Winarsih, S. (2021) Generating the responses
immune with honey, *Saussurea costus*, and *Nigella sativa* in cellular and humoral
may resolve COVID-19? *Syst. Rev. Pharm.*, 12(1): 1588-1593.
31. Schroeder, G.N. and Hilbi, H. (2008) Molecular pathogenesis of *Shigella* spp.:
Controlling host cell signaling, invasion, and death by Type III secretion. *Clin.*
Microbiol. Rev., 21(1): 134-156.
32. Killackey, S.A., Sorbara, M.T. and Girardin, S.E. (2016) Cellular aspects of *Shigella*
pathogenesis: Focus on the manipulation of host cell processes. *Front. Cell. Infect.*
Microbiol., 6(38): 1-11.
33. Salyers, A.A. and Whitt, D.D. (2002) Bacterial Pathogenesis: A Molecular Approach.
2nd ed. American Society for Microbiology Press, Washington DC.
34. Nezet, V. and Esko, J.D. (2009) Bacterial and Viral Infections, Mechanism of
Colonizations and Invasion, Essential of Glycology. 2nd ed. La-Jolla, California.
35. Ashida, H., Mimuro, H. and Sasakawa, C. (2015) *Shigella* manipulates host immune
responses by delivering effector proteins with specific roles. *Front. Immunol.*, 6(219):
1-12

36. Rasolofo-Razanamparany, V., Cassel-Beraud, A.M., Roux, J., Sansonetti, P.J. and Phalipon, A. (2001) predominance of serotype-specific mucosal antibody response in *Shigella flexneri*-infected humans living in an area of endemicity. *Infect. Immun.*, 69(9): 5230-5234.
37. Leung, S., Liu, X., Fang, L., Chen, X., Guo, T. and Zhang, J. (2010) The cytokine milieu in the interplay of pathogenic Th1/Th17 cells and regulatory T cells in autoimmune disease. *Cell. Mol. Immunol.*, 7(3): 182-189.
38. Abbas, A.K., Lichtman, A.H. and Pillai, S. (2015) Cellular and Molecular Immunology. 8th ed. Elsevier-Saunders, United States.
39. Kaetzel, C.S. (2007) Mucosal Immune Defense: Immunoglobulin A. Springer, New York.
40. Sperandio, B., Regnault, B., Guo, J., Zhang, Z., Stanley, S.L.J., Sansonetti, P.J. and Pédrón, T. (2008) Virulent *Shigella flexneri* subverts the host innate immune response through manipulation of antimicrobial peptide gene expression. *J. Exp. Med.*, 205(5): 1121-1132.
41. Dhanda, S.K., Gupta, S., Vir, P. and Raghava, G.P.S. (2013) Prediction of IL-4 Inducing Peptides. *Clin. Dev. Immunol.*, 2013(Article ID 263952): 1-9

42. Chitradevi, S.T.S., Kaur, G., Uppalapati, S., Yadav, A., Singh, D. and Bansal, A. (2015) Co-administration of rIpaB domain of *Shigella* with rGroEL of *S. Typhi* enhances the immune responses and protective efficacy against *Shigella* infection. *Cell. Mol. Immunol.*, 12(6): 757-767.
43. Lee, S.J., McLachlan, J.B., Kurtz, J.R., Fan, D., Winter, S.E., Bäumlner, A.J, Jenkins, M.K. and McSorley, S.J. (2012) Temporal expression of bacterial proteins instructs host CD4 T cell expansion and Th17 development. *PLoS Pathog.*, 8(1): e1002499.
44. Kurtz, J.R., Petersen, H.E., Frederick, D.R., Morici, L.A. and McLachlan, J.B. (2014) Vaccination with a single CD4 T cell peptide epitope from a *Salmonella* Type III-secreted effector protein provides protection against lethal infection. *Infect. Immun.*, 82(6): 2424-2433.
45. Brown, M.A. and Hural, J. (1997) Functions of IL-4 and control of its expression. *Crit. Rev. Immunol.*, 17(1): 1-32.
46. Padh, H., Yagnik, B., Sharma, D. and Desai, P. (2021) EpiMix based novel vaccine candidate for *Shigella*: Evidence of prophylactic immunity in Balb/c Mice. *Int. J. Pept. Res. Ther.*, 30 January): 1-16

47. León, Y., Zapata, L., Molina, R.E., Okanovic, G., Gómez, L.A., Daza-Castro, C., Flores-Concha, M., Reyes, J.L. and Oñate, A.A. (2020) Intranasal immunization of mice with multiepitope chimeric vaccine candidate based on conserved autotransporters SigA, Pic and Sap, confers protection against *Shigella flexneri*. *Vaccines*, 8(4): 563.
48. Jennison, A.V. and Verma, N.K. (2004) *Shigella flexneri* infection: Pathogenesis and vaccine development. *FEMS. Microbiol. Rev.*, 28(1): 43-58.
49. Pasetti, M.F., Venkatesan, M.M. and Barry, E.M. (2020) Chapter 30-oral *Shigella* vaccines. In: Kiyono, H. and Pascual, D.W., editors. *Mucosal Vaccines*. 2nd ed. Academic Press, Cambridge, MA, USA. p515-536.
50. Ishigame, H., Kakuta, S., Nagai, T., Kadoki, M., Nambu, A., Komiyama, Y., Fujikado, N., Tanahashi, Y., Akitsu, A., Kotaki, H., Sudo, K., Nakae, S., Sasakawa, C. and Iwakura, Y. (2009) Differential roles of interleukin-17A and -17F in host defense against mucosal bacterial infection and allergic responses. *Immunity*, 30(1): 108-119.
51. Curtis, M.M. and Way, S.S. (2009) Interleukin-17 in host defense against bacterial, mycobacterial and fungal pathogens. *Immunology*, 126(2): 177-185.

52. Martinez-Becerra, F.J., Scobey, M., Harrison, K., Choudhari, S.P., Quick, A.M., Joshi, S.B., Middaugh, C.R. and Picking, W.L. (2013) Parenteral immunization with IpaB/IpaD protects mice against lethal pulmonary infection by *Shigella*. *Vaccine*, 31(24): 2667-2672.
53. Blaschitz, C. and Raffatellu, M. (2010) Th 17 cytokines and the gut mucosal barrier. *J. Clin. Immunol.*, 30(2): 196-203.
54. Agerberth, B., Bergman, P. and Gudmundsson, G.H. (2013) Helping the host: Induction of antimicrobial peptides as a novel therapeutic strategy against infections. In: *Antimicrobial Peptides and Innate Immunity*. Springer, Basel, Switzerland. p359-375.
55. Cobo, E.R. and Chadee, K. (2013) Antimicrobial human β -defensins in the colon and their role in infectious and non- infectious diseases. *Pathogens*, 2(1): 177-192.
56. Fusco, A., Savio, V., Cammarota, M., Alfano, A., Schiraldi, C. and Donnarumma, G. (2017) Beta-defensin-2 and beta-defensin-3 reduce intestinal damage caused by *Salmonella typhimurium* modulating the expression of cytokines and enhancing the probiotic activity of *Enterococcus faecium*. *J. Immunol. Res.*, 2017;(Article ID 6976935): 1-9.

57. Wang, S., Huiqin L., Xinyi, Z. and Feng, Q. (2015) Intranasal and oral vaccination with protein-based antigens: Advantages, challenges, and formulation strategies.

Protein Cell, 6(7): 480-503.

58. Coria, L.M., Gabriela, S.R., Francisco, F.G., Mariana, C.F., Laura, B., Karina, A. and Cassataro, J. (2019) Oral co-administration of a bacterial protease inhibitor in the vaccine formulation increases antigen delivery at the intestinal epithelial barrier. *J. Control. Release.*, 293(???): 158-171.

Commented [TPS23]: Kindly provide issue number

Commented [Microsoft24R23]: this article did not provide the issue number





Figure-1: The secondary structure modeling of selected peptide. (a) pep_1 (QSSTGTNSQSDLDS) 14-mer. (b) pep_2 (DTTITKAETKTVTKNQVVDTPVTTDAAK) 28-mer and (c) pep_3 (ATLGATLNRLDFNVNKK) 17-mer.







Figure-2: Cellular immune response of intestinal mesenteric tissue of mice following oral immunization was determined by flow cytometry. (a) The significant increase of Th2 expressing CD4-IL4 is significantly higher in immunized groups (b). The significant increase of Th17 expressing CD4-IL17 is significantly higher in immunized groups, especially in CTB+pep_1 group. Data are shown as mean±SD.

???

Commented [TPS25]: Kindly provide the image for figure 3







Figure-3: Humoral immune response of intestinal mesenteric tissue of mice following oral immunization was determined by ELISA. (a) The significant increase of IL-4 in CTB_pep_2 and CTB_pep 3 groups (b). The significant increase of IL-17 in CTB+pep_1 group. Data are shown as mean±SD.






Figure-4: Cellular immune response of intestinal mucosal layer of mice following oral immunization was determined by ELISA. (a) The significant increase of sIgA in CTB_pep_3 group (b). The significant differences of β -defensin level among immunized groups. Data are shown as mean \pm SD.

